

ARTINIANNES OF LOCAL COHOMOLOGY MODULES OF ZD -MODULES

KAMRAN DIVAANI-AAZAR AND MOHAMMAD ALI ESMKHANI

ABSTRACT. This paper centers around Artinianness of the local cohomology of ZD -modules. Let $\mathfrak{a}$ be an ideal of a commutative Noetherian ring R . The notion of $\mathfrak{a}$ -relative Goldie dimension of an R -module M , as a generalization of that of Goldie dimension is presented. Let M be a ZD -module such that $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite. It is shown that if $\dim R/\mathfrak{a} = 0$, then the local cohomology modules $H_{\mathfrak{a}}^i(M)$ are Artinian. Also, it is proved that if $d = \dim M$ is finite, then $H_{\mathfrak{a}}^d(M)$ is Artinian, for any ideal $\mathfrak{a}$ of R . These results extend the previously known results concerning Artinianness of local cohomology of finitely generated modules.

1. INTRODUCTION

Throughout this paper, R is a commutative Noetherian ring with identity and all modules are assumed to be unitary. Let $\mathfrak{a}$ be an ideal of R and M an R -module. The $\mathfrak{a}$ -torsion submodule $\cup_{n \in \mathbb{N}} (0 :_M \mathfrak{a}^n)$ of M is denoted by $\Gamma_{\mathfrak{a}}(M)$. For each integer $i \geq 0$, the i -th local cohomology functor $H_{\mathfrak{a}}^i(\cdot)$ is defined as the i -th right derived functor of $\mathfrak{a}$ -torsion functor $\Gamma_{\mathfrak{a}}(\cdot)$. Also, it is known that for each $i \geq 0$ there is a natural isomorphism of R -modules

$$H_{\mathfrak{a}}^i(M) \cong \varinjlim_n \text{Ext}_R^i(R/\mathfrak{a}^n, M).$$

We refer the reader to text book [2] for more details about local cohomology.

It is known that the local cohomology of finitely generated modules have many interesting properties. In particular, if $(R, \mathfrak{m})$ is a local ring and M a finitely generated R -module, then the local cohomology modules $H_{\mathfrak{m}}^i(M)$ are Artinian. Also, in the same situation, it is known that for $d = \dim M$, the d -local cohomology module of M with respect to any ideal $\mathfrak{a}$ is Artinian. It will be a noticeable achievement, if

2000 *Mathematics Subject Classification.* 13D45, 13E10.

Key words and phrases. Local cohomology, Artinian modules, ZD -modules, Goldie dimension. This work has been supported by the Research Institute for Fundamental Sciences, Tabriz, Iran.

we could extend these results to local cohomology of a larger class of modules. In this paper, we shall show that ZD -modules behave very well in conjunction with Artinianness of local cohomology modules.

An R -module M is said to be ZD -module (zero-divisor module) if for any submodule N of M , the set of zero divisors of M/N is a union of finitely many prime ideals in $\text{Ass}_R(M/N)$. According to Example 2.2, the class of ZD -modules is much larger than that of finitely generated modules. As the main result of this paper, we prove that for a ZD -module M the following are equivalent:

- i) $\Gamma_{\mathfrak{a}}(M/N)$ is Artinian for any submodule N of M .
- ii) $H_{\mathfrak{a}}^i(M/N)$ is Artinian for any submodule N of M and all $i \geq 0$.

We say that an R -module M has finite $\mathfrak{a}$ -relative Goldie dimension if the Goldie dimension of the $\mathfrak{a}$ -torsion submodule of M is finite. Clearly, $\mathfrak{a}$ -relative Goldie dimension of any finitely generated module is finite. Let M be a ZD -module such that $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite. By using the above mentioned result, we deduce that the local cohomology modules $H_{\mathfrak{a}}^i(M)$ are Artinian if either,

- i) $\dim R/\mathfrak{a} = 0$, or
- ii) $d = \dim M$ is finite and $i = d$.

2. ZD -MODULES AND GOLDIE DIMENSION

Let for an R -module M , $Z_R(M)$ denote the set of zero divisors on M . Evans [4] calls a ring R a ZD -ring (zero-divisor ring) if for any ideal $\mathfrak{a}$ of R , $Z_R(R/\mathfrak{a})$ is a union of finitely many prime ideals. Next, we present the following modification of the definition of ZD -modules in [6].

Definition 2.1. An R -module M is said to be ZD -module if for every submodule N of M , the set $Z_R(M/N)$ is a union of finitely many prime ideals in $\text{Ass}_R(M/N)$.

An R -module M is said to be Laskerian if any submodule of M is an intersection of a finite number of primary submodules. Obviously, any Noetherian module is Laskerian. An R -module M [3] is said to be weakly Laskerian if the set of associated primes of any quotient module of M is finite. Clearly, any Laskerian module is weakly Laskerian and any weakly Laskerian module is ZD -module. In the sequel, we provide a large variety of examples of ZD -modules.

Example 2.2. i) It is easy to see that any module with finite support is weakly Laskerian. In particular, any Artinian module is a ZD -module. Also, by using this fact we can provide examples of ZD -modules which are neither finitely generated nor Artinian.

ii) Recall that a module M is said to have finite Goldie dimension if M does not contain an infinite direct sum of non-zero submodules, or equivalently, the injective envelope $E(M)$ of M decomposes as a finite direct sum of indecomposable injective submodules. Because for any R -module C , we have $\text{Ass}_R C = \text{Ass}_R E(C)$, it turns out that any module with finite Goldie dimension has only finitely many associated prime ideals. This yields that a module of which all quotients have finite Goldie dimension is weakly Laskerian.

iii) Let E be the minimal injective cogenerator of R and M an R -module. If for an R -module M the natural map from M to $\text{Hom}_R(\text{Hom}_R(M, E), E)$ is an isomorphism, then M is said to be Matlis reflexive. By [1, Theorem 12], an R -module M is Matlis reflexive if and only if M has a finitely generated submodule S such that M/S is Artinian and $R/\text{Ann}_R M$ is a complete semi-local ring. Also, by [5, Corollary 1.2], any quotient of an R -module M has finite Goldie dimension if and only if M has a finitely generated submodule S such that M/S is Artinian. Thus, by (ii) any Matlis reflexive module is a ZD -module.

iv) An R -module M is said to be linearly compact if each system of congruences

$$x \equiv x_i(M_i),$$

indexed by a set I and where the M_i are submodules of M , has a solution x whenever it has a solution for every finite subsystem. It is clear that, every quotient of a linearly compact module is also linearly compact. On the other hand a linearly compact module M has finite Goldie dimension (see e.g. [9, Chapter 1.3]). Thus, if M is a linearly compact module, then any quotient of M has finite Goldie dimension, and so, by (ii) M is a ZD -module.

Next, we bring the following characterization of ZD -modules.

Lemma 2.3. *Let M be an R -module. The following are equivalent:*

- i) M is a ZD -module.
- ii) For every submodule N of M , the number of prime ideals with the property being maximal in $\text{Ass}_R(M/N)$ is finite.

Proof. The proof is easy and we left it to the reader. $\square$

Lemma 2.4. *Let $\mathfrak{a}$ be a non-zero ideal of R and M a ZD -module. If M is $\mathfrak{a}$ -torsion free, then $\mathfrak{a}$ contains a nonzero divisor on M .*

Proof. Since M is ZD -module, there are prime ideals $\mathfrak{p}_1, \mathfrak{p}_2, \dots, \mathfrak{p}_n$ in $\text{Ass}_R M$ such that $Z_R(M) = \cup_{i=1}^n \mathfrak{p}_i$. Because, M is $\mathfrak{a}$ -torsion free, it follows that $\mathfrak{a}$ is not contained in any associated prime ideal of M . Thus, by Prime Avoidance Theorem, $\mathfrak{a}$ is not contained in $Z_R(M)$. $\square$

For an R -module M , the Goldie dimension of M is defined as the cardinal of the set of indecomposable submodules of $E(M)$, which appear in a decomposition of $E(M)$ into direct sum of indecomposable submodules. We shall use $\text{Gdim } M$ to denote the Goldie dimension of M . For a prime ideal $\mathfrak{p}$, let $\mu^0(\mathfrak{p}, M)$ denote the 0-th Bass number of M with respect to prime ideal $\mathfrak{p}$. It is known that $\mu^0(\mathfrak{p}, M) > 0$ if and only if $\mathfrak{p} \in \text{Ass}_R M$. It is clear by the definition of Goldie dimension that $\text{Gdim } M = \sum_{\mathfrak{p} \in \text{Spec } R} \mu^0(\mathfrak{p}, M)$. Having this in mind, we introduce the following generalization of the notion of Goldie dimension.

Definition 2.5. Let $\mathfrak{a}$ be an ideal of R . For an R -module M , we define $\mathfrak{a}$ -relative Goldie dimension of M as $\text{Gdim}_{\mathfrak{a}} M := \sum_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M)$. Here $V(\mathfrak{a})$ denotes the set of prime ideals of R which are containing $\mathfrak{a}$.

Obviously, if $\mathfrak{a}$ is the zero ideal, then $\text{Gdim}_{\mathfrak{a}} M = \text{Gdim } M$. Also, it is clear that the Goldie dimension of any Noetherian module as well as any Artinian module is finite.

Lemma 2.6. *Let $\mathfrak{a}$ be an ideal of R and M an R -module. Then $\text{Gdim}_{\mathfrak{a}} M = \text{Gdim } \Gamma_{\mathfrak{a}}(M)$.*

Proof. Let $\mathfrak{p}$ be a prime ideal of R . By [7, Theorem 18.4], each element of $E(R/\mathfrak{p})$ is annihilated by some power of $\mathfrak{p}$ and for each element $r \in R \setminus \mathfrak{p}$, the multiplication by r induces an automorphism of $E(R/\mathfrak{p})$. Therefore, it follows that $E(R/\mathfrak{p})$ is $\mathfrak{a}$ -torsion if $\mathfrak{a} \subseteq \mathfrak{p}$, and $\mathfrak{a}$ -torsion free otherwise. Hence $\Gamma_{\mathfrak{a}}(E(M)) = \bigoplus_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M) E(R/\mathfrak{p})$. It is easy to see that $\Gamma_{\mathfrak{a}}(E(M))$ is an essential extension of $\Gamma_{\mathfrak{a}}(M)$. On the other hand $\Gamma_{\mathfrak{a}}(E(M))$ is an injective R -module by [2, Proposition 2.1.4]. Hence $\Gamma_{\mathfrak{a}}(E(M)) \cong E(\Gamma_{\mathfrak{a}}(M))$. Thus

$$\text{Gdim}_{\mathfrak{a}} M = \sum_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M) = \text{Gdim } \Gamma_{\mathfrak{a}}(M). \square$$

Lemma 2.7. *Let $\mathfrak{a}$ be an ideal of R and M a ZD -module. The following are equivalent:*

- i) $\text{Gdim}_{\mathfrak{a}} M$ is finite.*
- ii) $\text{Gdim}_{\mathfrak{a}R_{\mathfrak{p}}} M_{\mathfrak{p}}$ is finite for any prime ideal $\mathfrak{p}$ of R .*
- iii) $\text{Gdim}_{\mathfrak{a}R_{\mathfrak{p}}} M_{\mathfrak{p}}$ is finite for any prime ideal $\mathfrak{p}$ which is maximal in $\text{Ass}_R M$.*

Proof. First we show that (i) implies (ii). Let $\mathfrak{p}$ be a prime ideal of R and S a multiplicatively closed subset of R . It follows by [2, Lemma 10.1.12], that if $S \cap \mathfrak{p} = \emptyset$, then the $S^{-1}R$ -modules $S^{-1}(E(R/\mathfrak{p}))$ and $E(R/\mathfrak{p})$ are isomorphic. Also, if $S \cap \mathfrak{p} \neq \emptyset$ we can easily deduce that $S^{-1}(E(R/\mathfrak{p})) = 0$. Thus, we have

$$S^{-1}(E(\Gamma_{\mathfrak{a}}(M))) \cong \bigoplus_{\mathfrak{p} \in V(\mathfrak{a}), \mathfrak{p} \cap S = \emptyset} \mu^0(\mathfrak{p}, M) E(R/\mathfrak{p}).$$

On the other hand for any R -module N , it follows by [2, Corollary 11.1.6], that as an $S^{-1}R$ -module $S^{-1}(E(N))$ is isomorphic to the $E_{S^{-1}R}(S^{-1}N)$. Thus

$$E_{S^{-1}R}(\Gamma_{\mathfrak{a}S^{-1}R}(S^{-1}M)) \cong E_{S^{-1}R}(S^{-1}(\Gamma_{\mathfrak{a}}(M))) \cong \bigoplus_{\mathfrak{p} \in V(\mathfrak{a}), \mathfrak{p} \cap S = \emptyset} \mu^0(\mathfrak{p}, M) E_{S^{-1}R}(S^{-1}R/S^{-1}\mathfrak{p}).$$

This shows that $\text{Gdim}_{\mathfrak{a}S^{-1}R} S^{-1}M \leq \text{Gdim}_{\mathfrak{a}} M$. Therefore (i) implies (ii), as required.

Clearly, (ii) implies (iii). Next, we prove that (iii) implies (i). Let $\{\mathfrak{q}_1, \mathfrak{q}_2, \dots, \mathfrak{q}_n\}$ be the set of all prime ideals with the property being maximal in $\text{Ass}_R M$. Note that by Lemma 2.3, this set is finite. Fix $1 \leq i \leq n$. As shown in the proof of Lemma 2.6, we have $E(\Gamma_{\mathfrak{a}}(M)) = \bigoplus_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M) E(R/\mathfrak{p})$. Thus

$$\begin{aligned} E_{R_{\mathfrak{q}_i}}(\Gamma_{\mathfrak{a}R_{\mathfrak{q}_i}}(M_{\mathfrak{q}_i})) &\cong (E(\Gamma_{\mathfrak{a}}(M)))_{\mathfrak{q}_i} \\ &\cong (\bigoplus_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M) E(R/\mathfrak{p}))_{\mathfrak{q}_i} \\ &\cong \bigoplus_{\mathfrak{a} \subseteq \mathfrak{p} \subseteq \mathfrak{q}_i} \mu^0(\mathfrak{p}, M) E_{R_{\mathfrak{q}_i}}(R_{\mathfrak{q}_i}/\mathfrak{p}R_{\mathfrak{q}_i}). \end{aligned}$$

Hence we have $\text{Gdim}_{\mathfrak{a}R_{\mathfrak{q}_i}} M_{\mathfrak{q}_i} = \sum_{\mathfrak{a} \subseteq \mathfrak{p} \subseteq \mathfrak{q}_i} \mu^0(\mathfrak{p}, M)$, and so

$$\begin{aligned} \text{Gdim}_{\mathfrak{a}R_{\mathfrak{q}_i}} M_{\mathfrak{q}_i} &= \sum_{\mathfrak{a} \subseteq \mathfrak{p} \subseteq \mathfrak{q}_i} \mu^0(\mathfrak{p}, M) \\ &\leq \text{Gdim}_{\mathfrak{a}} M \\ &\leq \sum_{i=1}^n (\sum_{\mathfrak{a} \subseteq \mathfrak{p} \subseteq \mathfrak{q}_i} \mu^0(\mathfrak{p}, M)) \\ &= \sum_{i=1}^n (\text{Gdim}_{\mathfrak{a}R_{\mathfrak{q}_i}} M_{\mathfrak{q}_i}). \end{aligned}$$

This concludes the proof. $\square$

3. ARTINIANNES OF LOCAL COHOMOLOGY MODULES

In [8, Theorem 1.3], Melkersson proved that an $\mathfrak{a}$ -torsion module M is Artinian if and only if $0 :_M \mathfrak{a}$ is Artinian. In this section, we use this result to deduce several results concerning Artinianness of local cohomology of ZD -modules.

Theorem 3.1. *Let $\mathfrak{a}$ be a non-zero ideal of R and M a ZD -module. The following are equivalent:*

- i) $\Gamma_{\mathfrak{a}}(M/N)$ is Artinian for any submodule N of M .
- ii) $H_{\mathfrak{a}}^i(M/N)$ is Artinian for any submodule N of M and all $i \geq 0$.

Proof. It is clear that (ii) implies (i). Next, we show that (i) implies (ii) by using induction on i . The claim for $i = 0$ holds by the assumption. Assume that $i > 0$ and that the assertion holds for $i - 1$. Thus $H_{\mathfrak{a}}^{i-1}(M/N)$ is Artinian for all submodules N of M . Let N be a submodule of M and $X = M/N$. Because $H_{\mathfrak{a}}^i(X) \cong H_{\mathfrak{a}}^i(X/\Gamma_{\mathfrak{a}}(X))$, we may assume that X is $\mathfrak{a}$ -torsion free. Note that any quotient of a ZD -module is also a ZD -module. Since X is $\mathfrak{a}$ -torsion free, by Lemma 2.4, it follows that $\mathfrak{a}$ contains an element r which is nonzero divisor on X . The exact sequence

$$0 \longrightarrow X \xrightarrow{r} X \longrightarrow X/rX \longrightarrow 0,$$

induces an exact sequence

$$H_{\mathfrak{a}}^{i-1}(X/rX) \longrightarrow H_{\mathfrak{a}}^i(X) \xrightarrow{r} H_{\mathfrak{a}}^i(X).$$

By inductive hypothesis $H_{\mathfrak{a}}^{i-1}(X/rX)$ is Artinian, so that by using the above exact sequence, we deduce that $(0 :_{H_{\mathfrak{a}}^i(X)} r)$ is Artinian. Since $H_{\mathfrak{a}}^i(X)$ is Rr -torsion, the conclusion follows by [8, Theorem 1.3]. $\square$

When $\dim R/\mathfrak{a} = 0$, we may strengthen Theorem 3.1 as follows.

Proposition 3.2. *Let the situation be as in Theorem 3.1. In addition assume that $\dim R/\mathfrak{a} = 0$. The following are equivalent:*

- i) $\Gamma_{\mathfrak{a}}(M/N)$ is Artinian for any submodule N of M .
- ii) $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite.
- iii) $H_{\mathfrak{a}}^i(M/N)$ is Artinian for any submodule N of M and all $i \geq 0$.
- iv) $H_{\mathfrak{a}R_{\mathfrak{p}}}^i(M_{\mathfrak{p}}/N_{\mathfrak{p}})$ is Artinian for any submodule N of M , any prime ideal $\mathfrak{p}$ of R and all $i \geq 0$.

Proof. In view of Theorem 3.1, Lemma 2.6 and Lemma 2.7, it suffices to show that an $\mathfrak{a}$ -torsion module M is Artinian if and only if its Goldie dimension is finite. Assume that M is a $\mathfrak{a}$ -torsion module. Then $\text{Ass}_R M \subseteq V(\mathfrak{a})$. On the other hand, because $\dim R/\mathfrak{a} = 0$, it turns out that $V(\mathfrak{a})$ is a finite set consisting of maximal ideals. It is clear that if M is Artinian, then the Goldie dimension of M has to be finite. Conversely, suppose that the Goldie dimension of M is finite. Then

$$\sum_{\mathfrak{p} \in \text{Ass}_R M} \mu^0(\mathfrak{p}, M) \leq \sum_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M) < \infty.$$

Thus $E(M)$ is direct sum of finitely many Artinian modules. $\square$

Let $\mathfrak{a}$ be an ideal of a local ring $(R, \mathfrak{m})$ and let M be a finitely generated R -module of dimension d . By [2, Theorem 7.1.6], $H_{\mathfrak{a}}^d(M)$ is Artinian. Also, it is known that if $\dim R/\mathfrak{a} = 0$, then $H_{\mathfrak{a}}^i(M)$ is Artinian for all $i \geq 0$. Next, we provide a far reaching generalization of these facts.

Corollary 3.3. *Let $\mathfrak{a}$ be an ideal of R and M a ZD-module. Assume that $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite. We have the following.*

- i) *If $\dim R/\mathfrak{a} = 0$, then $H_{\mathfrak{a}}^i(M)$ is Artinian for all $i \geq 0$.*
- ii) *If $d = \dim M$ is finite, then $H_{\mathfrak{a}}^d(M)$ is Artinian.*

Proof. i) is clear by Proposition 3.2.

ii) We use induction on d . Suppose $d = 0$. Then every associated prime ideal of M is maximal and so $E(\Gamma_{\mathfrak{a}}(M))$ is a direct sum of a finitely many $E(R/\mathfrak{m})$, where $\mathfrak{m}$'s are maximal ideals of R . Hence $\Gamma_{\mathfrak{a}}(M)$ is Artinian.

Now, we assume that $d > 0$ and that the claim holds for $d - 1$. Similar to the proof of Theorem 3.1, we may assume that M is $\mathfrak{a}$ -torsion free. Thus we can choose an element $r \in \mathfrak{a}$, which is nonzero divisor on M . From the exact sequence

$$0 \longrightarrow M \xrightarrow{r} M \longrightarrow M/rM \longrightarrow 0,$$

we deduce the exact sequence

$$H_{\mathfrak{a}}^{d-1}(M/rM) \longrightarrow H_{\mathfrak{a}}^d(M) \xrightarrow{r} H_{\mathfrak{a}}^d(M),$$

of local cohomology modules. Since r is a nonzero divisor on M , we have $\dim M/rM \leq d - 1$. Hence, it follows from inductive hypothesis or Grothendieck's Vanishing Theorem [2, Theorem 6.1.2] that $H_{\mathfrak{a}}^{d-1}(M/rM)$ is Artinian. Therefore by using [8, Theorem 1.3], we deduce that $H_{\mathfrak{a}}^d(M)$ is Artinian. $\square$

Next, we bring an example to show that there is a non-finitely generated ZD -module M , and an ideal $\mathfrak{a}$ of R , such that $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite.

Example 3.4. i) Let M be a Matlis reflexive R -module and $\mathfrak{a}$ an arbitrary ideal of R . Then it follows, by Example 2.2(iii), that M is a ZD -module and that $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite.

ii) Let $\mathfrak{m}, \mathfrak{n}$ be two distinct maximal ideals of a ring R . Put $M = \bigoplus_{i \in \mathbb{N}} R/\mathfrak{m}$ and $\mathfrak{a} = \mathfrak{n}$. Then M is a ZD -module and $\mathfrak{a}$ -relative Goldie dimension of any quotient of M is finite. Also, note that by Example 2.2(iii), M is not Matlis reflexive.

REFERENCES

- [1] R. Belshoff, E. Enochs and J.R. Garcia Rozas, *Generalized Matlis duality*, Proc. Amer. Math. Soc., **128**(5) (2000), 1307-1312.
- [2] M.P. Brodmann and R.Y. Sharp: '*Local cohomology-An algebraic introduction with geometric applications*', Cambr. Univ. Press, 1998.
- [3] K. Divaani-Aazar and A. Mafi, *Associated primes of local cohomology modules*, Proc. Amer. Math. Soc., to appear.
- [4] E.G. Evans, *Zero divisors in noetherian-like rings*, Trans. Amer. Math. Soc., **155** (1971), 505-512.
- [5] C. Faith and D. Herbera, *Endomorphism rings and tensor products of linearly compact modules*, Comm. Algebra, **25**(4) (1997), 1215-1255.
- [6] W. Heinzer and D. Lantz, *The Laskerian property in commutative rings*, J. Algebra **72**(1) (1981), 101-114.
- [7] H. Matsumura, *Commutative ring theory*, Second edition, Cambridge Studies in Advanced Mathematics, **8**, Cambridge University Press, Cambridge, 1989.
- [8] L. Melkersson, *On asymptotic stability of prime ideals connected with the powers of an ideal*, Math. Proc. Camb. Phil. Soc., **107** (1990), 267-271.
- [9] W. Xue, *Rings with Morita duality*, Lecture Notes in Mathematics, **1523**, Springer-Verlag, Berlin, 1992.

K. DIVAANI-AAZAR, DEPARTMENT OF MATHEMATICS, AZ-ZAHRA UNIVERSITY, VANAK, POST CODE 19834, TEHRAN, IRAN-AND-RESEARCH INSTITUTE FOR FUNDAMENTAL SCIENCES, TABRIZ, IRAN.

E-mail address: kdivaani@ipm.ir

M.A. ESMKHANI, DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, TEHRAN, IRAN.